


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 7/48, C07D 405/06	A1	(11) International Publication Number: WO 93/18743 (43) International Publication Date: 30 September 1993 (30.09.93)
(21) International Application Number: PCT/EP93/00599 (22) International Filing Date: 12 March 1993 (12.03.93) (30) Priority data: 92200797.6 20 March 1992 (20.03.92) DE (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). (72) Inventor; and (75) Inventor/Applicant (for US only) : CAUWENBERGH, Gerard, Frans, Maria, Jan [BE/BE]; Tuinweg 38, B-2290 Vorselaar (BE). (74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica N.V., Patent Department, Turnhoutseweg 30, B-2340-Berse (BE).		(81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: AGENT FOR REGULATING THE GREASINESS OF THE SKIN**(57) Abstract**

The use of a compound having formula ((cis)-(I)), wherein R represents methyl or ethyl, for regulating the greasiness of the skin, particular novel physical forms of said compound, cosmetic compositions comprising said agent, processes for preparing said agent and compositions, and a method of reducing the greasiness of the skin.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

Agent for regulating the greasiness of the skin

10 A valued aspect of man is the appearance of his skin. A greasy skin, e.g. after a busy day or following physical exercise, is often considered an annoyance, even though it is temporary and does not indicate any health problem. The present invention concerns the use of an agent for regulating the greasiness of the skin, particular novel physical forms of said agent, cosmetic compositions comprising said agent, processes for preparing said agent and compositions, and a method of regulating the greasiness of the skin.

The present invention is concerned with the use of a compound having the formula



20

wherein R represents methyl or ethyl, preferably ethyl, or a cosmetically acceptable acid addition salt thereof, for regulating the greasiness of the skin of human beings.

25 The present invention is also concerned with a method of regulating the greasiness of the skin of human beings, which comprises administering topically to the skin of said human beings a compound having the formula (cis)-(I), or a cosmetically acceptable acid addition salt thereof, in an amount effective in regulating the greasiness of the skin.

30 The compound (I) can be applied to the skin when necessary or convenient, e.g. at each washing occasion or thereafter. The topical administration may be repeated until a cosmetically beneficial reduction of the greasiness of the skin is obtained. No special precautions are needed other than those which normally apply when using cosmetic agents.

The compounds of formula (I), (\pm)-methyl- or ethyl-cis-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinecarboxylate used in the method of the present invention are known agents and
5 their preparation as well as their properties are described in US-4,335,125.


The compound of formula (I) is preferably used as such or in a cosmetically acceptable acid addition salt form. Said salt forms can conveniently be prepared by treating the base form with an appropriate acid such as, for example, hydrohalic acids,
10 e.g. hydrochloric or hydrobromic acid; sulfuric acid; nitric acid; phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxy-propanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic,
15 benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids.

The term acid addition salt form as used hereinabove also comprises the solvates which the compound compound (I) and its acid addition salts are able to form. Examples of
20 such solvates are, e.g. the hydrates, alcoholates and the like.

The compound (I) is keratinophilic and binds to the keratin in the upper layers of the epidermis. When used in cosmetic compositions at concentrations of 0.05% to 1% (by weight based on the total weight of said cosmetic compositions), in particular 0.1% to 0.5% and preferably 0.1% to 0.2%, compound (I) exerts an inhibitory effect on the
25 formation of sterols such as cholesterol in the skin. Since this influences the lipid contents of the epidermis, a beneficial alteration of the greasiness of the skin results from administering compound (I) to the skin. In particular it appears that the biosynthesis of some free fatty acids is reduced and that the composition of said free fatty acids is altered. Compound (I) also affects the concentration of the phospholipids in the skin.
30 Besides affecting the lipid contents, the compound of formula (I) at the concentrations used also exerts some inhibitory effects on the growth of microorganisms such as bacteria and fungi.

Since said microorganisms depend on lipids occurring in the skin, the dual action of compound (I) on both the lipid contents of the skin and on the growth of
35 microorganisms ensures that the skin flora is checked and no unpleasant smells develop.

The compound (I) can be prepared as described in US-4,335,125 or preferably can be obtained from cis-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine (II) which is a commercial agent. In particular, compound (I) can be prepared by hydrolyzing cis-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine (II) and reacting the thus obtained product with a reagent of formula W-C(=O)-O-R (III) wherein R represents methyl or ethyl and W represents a leaving group such as halo, preferably chloro, methoxy or ethoxy.



Said hydrolysis can conveniently be conducted by stirring and heating a solution of cis-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine (II) in water in the presence of a base or an acid, preferably in the presence of concentrated hydrochloric acid. The hydrolysis product is extracted from the reaction mixture (neutralized with a base such as sodium or potassium hydroxide) with an organic solvent such as a halogenated hydrocarbon, e.g. dichloromethane, or an alcohol, e.g. methanol, ethanol and the like, or a mixture of such solvents. After evaporation of the organic layer, the crude hydrolysis product need not be purified and can be converted into compound (I) directly by dissolving it into a halogenated hydrocarbon, e.g. dichloromethane and treating it with methyl or ethyl chloroformate in the presence of a base such as sodium or potassium carbonate. Stirring of the reaction mixture enhances the reaction rate and the reaction can conveniently be conducted at room temperature.

In the method according to the present invention it is advantageous to use a micronized form of the compound (I), in particular material having an average particle size of less than 100 microns, preferably less than 75 microns, and in particular having a mean particle size of not more than 15 microns.

The micronized form of the compound (I) has the advantage of dissolving better and more rapidly due to its high surface area, and of penetrating well into the upper layers of the epidermis. Micronized forms of the compound (I) are novel and can be prepared by

micronization techniques known in the art, e.g. by milling in appropriate mills and sieving through appropriate sieves.

5 The free base form of compound (I) wherein R represents ethyl occurs in two polymorphic forms; the higher melting polymorph I as a substantially crystallographically pure substance can be characterized by its peak maximum at about 127.6°C with DSC; the lower melting polymorph II as a substantially crystallographically pure substance by its peak maximum at about 110.9°C. Polymorph I is more stable and slightly less soluble than polymorph II, and is the polymorphic form
10 used preferably in the method according to the present invention.

Polymorph I can conveniently be prepared by recrystallizing the free base form of compound (I) in a suitable solvent such as a ketone or an ester, e.g. acetone or ethyl acetate. Said recrystallization generally comprises dissolving the free base form of the compound of formula (I) in a hot solvent, concentrating the thus obtained solution,
15 allowing the concentrate to cool and separating the precipitated crystals. Preferably, the solution is treated with active charcoal before concentrating. Further, it may be advantageous to seed the supersaturated solution from which polymorph I is to be crystallized with appropriate crystals of said polymorph I that have been obtained in previous runs. Polymorph I can be micronized as described hereinbefore, by milling in
20 an appropriate mill and sieving the thus obtained product, for example, with an air-jet sifter using appropriate sieves such as 100 and 74 micrometer sieves.

Polymorph II on the other hand can be obtained by recrystallizing the free base form of compound (I) in a suitable solvent such as 2-propanol, hexane, or methylbenzene. Selective crystallization of said polymorph II may be induced by seeding the
25 supersaturated solution from which polymorph II is to be crystallized. Polymorph II also may be micronized although due care should be taken to avoid transition of polymorph II into polymorph I.

The compound (I) is most preferably applied to the affected areas of the body in the
30 form of appropriate compositions, in particular cosmetic compositions. Said compositions contain the compound (I), preferably in a 0.05 to 1% concentration (weight by weight), in particular in a 0.1% to 0.5% concentration and preferably in a 0.1% to 0.2% concentration, and any known dermatologically acceptable carrier. Said compositions may take a wide variety of forms such as, for example, liquid forms, e.g.
35 solutions, emulsions, gels or suspensions in aqueous, alcoholic or oily mediums, such as toilet waters, packs, lotions, skin milks or milky lotions and shampoos; or semi-liquid

formulations, e.g. creams, hydrogels, gels, pastes, ointments, salves, tinctures and the like, or solid formulations, e.g. powders.

Said preparations contain, besides the compound (I), components usually employed in such preparations. Examples of such components are oils, fats, waxes, surfactants, humectants, penetration enhancing agents, thickening agents, lipid absorbents, anti-oxidants, viscosity stabilizers, chelating agents, buffers, preservatives, perfumes, fragrances, dyestuffs, lower alkanols, unsaturated alcohols and the like. If desired, further active ingredients may be incorporated in the compositions, e.g. antiinflammatory agents, antibacterials, antifungals, woundhealing agents, disinfectants, vitamins, sunscreens, antibiotics or anti-dandruff agents.

The compound (I) is advantageously used in cosmetical products useful for dealing with an impure skin or for preventing the development of skin impurities such as blemishes, comedones, pimples and the like. Examples of such cosmetic compositions are sun protecting products to be used before, during or after exposure to the sun, products for tanning without sun, skin-whitening products, anti-wrinkle products, daily used hygienic preparations for cleaning and caring such as soaps, shampoos, bath and shower products, shaving creams, foams or lotions, baby care products, hygienic tissues, products for intimate hygiene care, e.g. intimate cleansing products, and daily used cosmetic preparations such as creams, emulsions, gels, oils, lotions, skin milks, milk gels, tinted milk gels, face masks, deodorants, anti-perspirants, perfumes, toilet waters, eau de Cologne, depilatories, after-shaves, make-up articles, e.g. creams, tinted bases, powders, lipstick, cover sticks, eye-liners; products for removing make-up from the face and the eyes; hair care products, e.g. hair tints and bleaches, products for waving, straightening and fixing, setting products, cleansing products, conditioning products or hairdressing products; and the like cosmetical products.

The liquid formulations may be packaged advantageously in any dispensing device adapted for topical administration, for example in flacons, bottles or also as a spray, either using an inert compressed gas as a propellant such as nitrogen or carbon dioxide, or alternatively using a pump to provide an aerosol. Solid formulations can be applied to the skin with powder puffs or directly with a cover stick. Alternatively, solid formulations such as granules, tablets or powders may also be dissolved in baths. Semi-liquid formulations can be packaged in suitable, art-known containers such as plastic, glass or ceramic pots, tubes, e.g. PVC-covered aluminum tubes.

Other particular compositions are those wherein the compound (I) is formulated in liposome-containing compositions. Different types of liposomes may be employed such as coarse (multilayer) liposomes or unilamellar liposomes and the like, which are formed, for example, with phosphatidyl cholines, ethanolamines, serines, sphingomyelins, cardiolipins, plasmalogens, phosphatidic acids, cerebroside and the like. The viscosity of the liposomes can be increased by addition of one or more thickening agents such as xanthan gum, hydroxypropyl cellulose, hydroxypropyl methylcellulose and mixtures thereof. The aqueous component may consist of water optionally in admixture with electrolytes, buffers and other ingredients such as preservatives. Preferred electrolytes are calcium, sodium and potassium chloride. The organic component may consist of a solvent such as ethanol, glycerol, propylene glycol, a polyethylene glycol and a suitable phospholipid such as, lecithin, phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl serine, phosphatidyl inositol, lysophosphatidyl choline, phosphatidyl glycerol and the like. Other lipophilic additives which may be added to selectively modify the characteristics of the liposomes are, e.g. stearylamine, phosphatidic acid, tocopherol, cholesterol, lanolin and the like.

For preparing ointments, creams, toilet waters, skin milks, and the like, compound (I) is combined in intimate admixture with a cosmetically acceptable carrier. In ointments or creams, the carrier for example consists of 1 to 20%, in particular 5 to 15% of a humectant, 0.1 to 10%, in particular from 0.5 to 5% of a thickener, and water; or said carrier may consist of 70 to 99%, in particular 80 to 95% of a surfactant, and 0 to 20%, in particular 2.5 to 15% of a fat; or 80 to 99.9%, in particular 90 to 99% of a thickener; or 5 to 15% of a surfactant, 2-15% of a humectant, 0 to 80% of an oil, small (0 to $\leq 2\%$) amounts of preservative, colouring agent and/or perfume, and water. In a toilet water, the carrier for example consists of 2 to 10% of a lower alcohol, 0.1 to 10% or in particular 0.5 to 1% of a surfactant, 1 to 20%, in particular 3 to 7% of a humectant, 0 to 5% of a buffer, water and small amounts (0 to $\leq 2\%$) of preservative, dyestuff and/or perfume. In a milk gel, the carrier typically consists of 50 to 80% of water, preferably 70 to 80% of water; 8 to 20%, preferably 10 to 15% of an alcohol; 1 to 10%, preferably 2 to 5% of a surfactant; 0.5 to 2% of one or more thickeners; and 0 to 3% of preservatives, humectants, nutritional factors, vitamins, astringents and/or perfumes. Other active ingredients may be incorporated at doses ranging from 0.005% to 0.5%, particularly from 0.01% to 0.1%.

Typical milk gel formulations comprise the active ingredient at the concentrations mentioned hereinabove; water; one or more gelling agents or thickeners such as, for example, cellulose derivatives, e.g. hydroxypropyl cellulose, hydroxypropyl

5 methylcellulose, or carboxyvinylpolymers at a concentration of 0.02% to 4%, particularly 0.2% to 2% and preferably about 0.2% to 1.5%; one or more alcohols such as, for example, ethanol, propanol, butanol, pentanol, glycerol, propylene glycol, dodecatrienol in a concentration of 0.02% to 50%, particularly 0.5% to 30% and
10 preferably about 0.5% to 20%. Said typical milk gel formulation may further comprise plant extracts, e.g. from aloe, arnica, birch, bladder wrack, gentian, ginseng, hamamelis (witch hazel), hawthorn, kina, lemon, nasturtium, rosemary and the like, at a concentration of from 0.1% to 35%, preferably from 0.5% to 12% and in particular from 1% to 3%; fatty ingredients such as paraffins at a concentration of from 0.1% to 30%,
15 preferably from 0.2% to 10% and in particular from 1% to 5%; vitamins such as, for example, vitamin E (tocopherol) and derivatives, e.g. tocopheryl acetate, panthenol, and the like, at a concentration of 0.01% to 10%, preferably 0.1% to 8% and in particular 0.5% to 2%; antiinflammatory products of synthetic or natural origin, e.g. bisabolol, at a concentration of 0% to 5%, preferably 0.1% to 3% and in particular 0.2% to 0.5%;
frangrances at a concentration of 0% to 2%, preferably 0.1% to 1% and in particular 0.3% to 0.5%; and optionally preservatives and/or antioxidants in an amount sufficient to prevent the degradation of the final composition.

Typical tinted formulations may be emulsions comprising the active ingredient at the concentrations mentioned hereinabove; water; one or more emulsifiers such as
20 ethoxylated fatty alcohols, e.g. ceteth, laureth, myreth, oleth, steareth, trideth at a concentration of 0.1% to 10%, preferably 0.5% to 5%; one or more fatty components such as a C₈₋₂₂ fatty alcohol, preferably a C₁₂₋₁₈ fatty alcohol at a concentration of 0.1% to 10%, preferably 0.5% to 5%, or a mineral oil (petrolatum) at a concentration of 0.1% to 80%, preferably 1% to 30%; or a polyoxyethylated or -propylated fatty alcohol ether,
25 e.g. polyoxypropylated stearyl ether, at a concentration of 1 to 40%, preferably 5 to 30%, and in particular 10 to 20%; one or more alcohols such as, for example, ethanol, propanol, butanol, pentanol, glycerol, propylene glycol, butylene glycol, dodecatrienol in a concentration of 0.1% to 25%, preferably 0.5% to 20%. Said tinted formulations obviously further comprise one or more pigments such as zinc oxide,
30 kaolin, mica, iron oxide, titaniumdioxide and the like. Said tinted formulations may further also comprise plant extracts, e.g. from aloe, arnica, birch, bladder wrack, gentian, ginseng, hamamelis (witch hazel), hawthorn, kina, lemon, nasturtium, rosemary and the like, at a concentration of from 0.1% to 35%, preferably from 0.5% to 12% and in particular from 1% to 3%; a gelling agent such as a cellulose derivative or a
35 carboxyvinyl polymer at a concentration of 0.1% to 1%; vitamins such as, for example, vitamin E (tocopherol) and derivatives, e.g. tocopheryl acetate, panthenol, and the like, at a concentration of 0.01% to 10%, preferably 0.1% to 8% and in particular 0.5% to

2%; antiinflammatory products of synthetic or natural origin, e.g. bisabolol, at a concentration of 0% to 5%, preferably 0.1% to 3% and in particular 0.2% to 0.5%; thickening agents, e.g. xanthan gum, at a concentration of 0.1% to 1%, preferably 0.2% to 0.5%; fragrances at a concentration of 0% to 2%, preferably 0.1% to 1% and in particular 0.3% to 0.5%; and optionally preservatives and/or antioxidants in an amount sufficient to prevent the degradation of the final composition. Preferably the above formulations have a pH ranging from 5 to 7.5, particularly from 5.5 to 7. Said pH can be established by the addition of a base, e.g. sodium hydroxide, or an acid, e.g. lactic, citric or phosphoric acid, or a buffer, e.g. a citrate, phosphate, lactate or acetate buffer.

10 A typical coverstick formulation comprises the active ingredient at the concentrations mentioned hereinabove; one or more fatty substances such as a mineral oil (petrolatum) at a concentration of 1% to 70%, preferably 30% to 60%, or a paraffin at a concentration of 0.1% to 30%, preferably 5% to 20%; triglycerides at a concentration of 0.1% to 30%, preferably 1% to 10%; a fatty C₈₋₂₂ alcohol, preferably a C₁₂₋₁₈ fatty alcohol at a concentration of 0.1% to 40%, preferably 1% to 10%; a wax at a concentration of 0.1% to 10%; and optionally a sufficient amount of preservatives and/or antioxidants to prevent the degradation of the final composition; and optionally other ingredients such as fragrances, woundhealing products, pigments and the like.

20 In the aforementioned preparations, all % symbols refer to weight by weight percentage. The humectant, surfactant, oil, other active ingredient, etc. referred to in said preparations may be any such component used in the cosmetic arts. Further, when in the above compositions one or more of the components make up the major part of the composition, the other ingredients can evidently be not present at their indicated maximum concentration and therefore will make up the remainder of the composition.

25 Experimental part

Example 1 : Preparation of compound (I)

30 a) A stirred mixture of 50 grams of cis-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-yl)methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine, 105 grams of water and 35 grams of concentrated hydrochloric acid was heated at 80-90°C for 16 hours. The reaction mixture was cooled to room temperature (approximately 20-25°C) and 70 ml of dichloromethane and 30 ml of methanol were added. There was added dropwise 25 ml of sodium hydroxide solution (50% in water) while stirring and keeping the temperature of the reaction mixture below 30°C by cooling with ice-water. After stirring for half an hour, there were added another 130 ml of dichloromethane and 170 ml of methanol. Stirring was continued for yet another half hour, the organic layer was separated and the

organic solvent was distilled off under reduced pressure. The residue was taken up in 200 ml of dichloromethane.

- b) To the residue there were added 52 grams of potassium carbonate and 12.25 grams of ethyl chloroformate were added dropwise while keeping the reaction temperature below 25°C. After stirring for two hours at room temperature, 112 grams of water were added and stirring was continued for four more hours. The organic layer was separated, dried, filtered and evaporated under reduced pressure. The residue was dissolved in 60 ml 4-methyl-2-pentanone by heating and crystallized by allowing the reaction mixture to cool spontaneously to room temperature. After cooling another four hours with ice-water, the precipitate was filtered off and dried in vacuo at 40°C, yielding 44.84 grams (84.9%) of (±)-ethyl cis-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinecarboxylate.
- c) A mixture of 6.5 grams of (±)-ethyl cis-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinecarboxylate in 51.5 ml of acetone was heated to the reflux temperature and 3.23 grams of active coal (Norit A supra) were added. After refluxing another half hour, the reaction mixture was filtered over diatomaceous earth while hot. The filter was washed with hot acetone and the combined filtrates were concentrated to an end volume of about 25 ml. The reaction mixture was allowed to cool to 40°C and was then seeded with polymorph I. The reaction mixture was allowed to cool further to room temperature and was then stirred for one hour while cooling with ice-water. The precipitate was filtered off and dried in vacuo at 40°C, yielding 5.35 parts (82.1%) of (±)-ethyl cis-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinecarboxylate (polymorph I); mp. 127.6°C.

Composition Examples

The following formulations exemplify typical compositions in accordance with the present invention.

- "Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or a cosmetically acceptable acid addition salt thereof.

Example 2 : 0.1% cream

- 75 mg stearyl alcohol, 2 mg cetyl alcohol, 20 mg sorbitan monostearate and 10 mg isopropyl myristate are introduced into a doublewall jacketed vessel and heated until the mixture has completely molten. This mixture is added to a separately prepared mixture of purified water, 200 mg propylene glycol and 15 mg polysorbate 60 having a temperature of 70 to 75°C while using a homogenizer for liquids. The resulting emulsion is allowed

to cool to below 25°C while continuously mixing. A solution of 1 mg A.I., 1 mg polysorbate 80 and purified water and a solution of 2 mg sodium sulfite anhydrous in purified water are next added to the emulsion while continuously mixing. The cream is homogenized and filled into suitable tubes.

5

Example 3 : 0.2% liposome formulation


A mixture of 0.2 g A.I. microfine, 20 g phosphatidyl choline, 5 g cholesterol and 10 g ethyl alcohol is stirred and heated at 55-60°C until complete dissolution and is added to a solution of 0.2 g methyl paraben, 0.02 g propyl paraben, 0.15 g disodium edetate and 10 0.3 g sodium chloride in purified water while homogenizing. 0.15 g Hydroxypropyl-methylcellulose in purified water ad 100 g is added and the mixing is continued until swelling is complete.

Example 4 : 0.1% liposome formulation

15 A mixture of 10 g phosphatidyl choline and 1 g cholesterol in 7.5 g ethyl alcohol is stirred and heated at 40°C until complete dissolution. 0.2 g A.I. microfine is dissolved in purified water by mixing while heating at 40°C. The alcoholic solution is added slowly to the aqueous solution while homogenizing during 10 minutes. 1.5 g Hydroxypropyl-methylcellulose in purified water is added while mixing until swelling is complete. The 20 resulting solution is adjusted to pH 5.0 with sodium hydroxide 1 N and diluted with the rest of the purified water ad 100 g.


Claims

1. The use of a compound having the formula



wherein R represents methyl or ethyl, or a cosmetically acceptable acid addition salt thereof, for regulating the greasiness of the skin.

- 10 2. A compound having the formula



15 wherein R represents methyl or ethyl, or a cosmetically acceptable acid addition salt thereof, characterized in that said compound is micronized and has a particle size of less than 100 micron.

- 20 3. A compound according to claim 2 wherein R represents ethyl and the compound is the higher melting polymorph of the free base form as a substantially crystallographically pure substance.
4. A process for the preparation of a micronized compound of formula (I) as defined in claim 2, characterized by milling and sieving said compound.
- 25 5. A process according to claim 4 for preparing the higher melting polymorph of the compound of formula (I) wherein R represents ethyl, characterized by recrystallizing the free base form of said compound in a ketone or ester as solvent, and milling and sieving the recrystallized compound.

6. A cosmetic composition for reducing the greasiness of the skin, comprising a cosmetic vehicle and as active ingredient a compound of formula (I) as defined in claims 2 or 3.
- 5 7. A cosmetic composition according to claim 6 comprising 0.05% to 1% of the active ingredient by weight based on the total weight of the cosmetic composition.
8. A cosmetic composition according to claim 7 comprising 0.1% to 0.2% of the active
10 ingredient by weight based on the total weight of the cosmetic composition.
9. A process of preparing a cosmetic composition as defined in any one of claims 6 to 8, characterized in that the micronized form of the compound of formula (I) is intimately mixed with a cosmetic vehicle.

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 A61K7/48; C07D405/06														
II. FIELDS SEARCHED <div style="text-align: center; margin-top: 10px;">Minimum Documentation Searched⁷</div> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%; padding: 5px;">Classification System</td> <td style="padding: 5px;">Classification Symbols</td> </tr> <tr> <td style="padding: 5px;">Int.Cl. 5</td> <td style="padding: 5px;">A61K ; C07D</td> </tr> </table> <div style="text-align: center; margin-top: 10px;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched⁸</div>			Classification System	Classification Symbols	Int.Cl. 5	A61K ; C07D								
Classification System	Classification Symbols													
Int.Cl. 5	A61K ; C07D													
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹ <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; padding: 5px;">Category¹⁰</th> <th style="width: 70%; padding: 5px;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 20%; padding: 5px;">Relevant to Claim No.¹³</th> </tr> <tr> <td style="padding: 5px; vertical-align: top;">A</td> <td style="padding: 5px;"> US,A,4 335 125 (HEERES ET AL.) 15 June 1982 cited in the application --- </td> <td style="padding: 5px;"></td> </tr> <tr> <td style="padding: 5px; vertical-align: top;">A</td> <td style="padding: 5px;"> WO,A,9 102 515 (HENKEL KOMMANDITGESELLSCHAFT AUF AKTIEN) 7 March 1991 --- </td> <td style="padding: 5px;"></td> </tr> <tr> <td style="padding: 5px; vertical-align: top;">P,A</td> <td style="padding: 5px;"> EP,A,0 490 583 (UNILEVER PLC) 17 June 1992 ----- </td> <td style="padding: 5px;"></td> </tr> </table> <div style="margin-top: 10px;"> <div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>¹⁰ Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div> </div>			Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	A	US,A,4 335 125 (HEERES ET AL.) 15 June 1982 cited in the application ---		A	WO,A,9 102 515 (HENKEL KOMMANDITGESELLSCHAFT AUF AKTIEN) 7 March 1991 ---		P,A	EP,A,0 490 583 (UNILEVER PLC) 17 June 1992 -----	
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³												
A	US,A,4 335 125 (HEERES ET AL.) 15 June 1982 cited in the application ---													
A	WO,A,9 102 515 (HENKEL KOMMANDITGESELLSCHAFT AUF AKTIEN) 7 March 1991 ---													
P,A	EP,A,0 490 583 (UNILEVER PLC) 17 June 1992 -----													
IV. CERTIFICATION <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;"> Date of the Actual Completion of the International Search <div style="text-align: center; margin-top: 10px;">18 MAY 1993</div> </td> <td style="width: 50%; padding: 5px;"> Date of Mailing of this International Search Report <div style="text-align: center; margin-top: 10px;">17. 06. 93</div> </td> </tr> <tr> <td style="width: 50%; padding: 5px;"> International Searching Authority <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div> </td> <td style="width: 50%; padding: 5px;"> Signature of Authorized Officer <div style="text-align: center; margin-top: 10px;">CHOULY J.</div> </td> </tr> </table>			Date of the Actual Completion of the International Search <div style="text-align: center; margin-top: 10px;">18 MAY 1993</div>	Date of Mailing of this International Search Report <div style="text-align: center; margin-top: 10px;">17. 06. 93</div>	International Searching Authority <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center; margin-top: 10px;">CHOULY J.</div>								
Date of the Actual Completion of the International Search <div style="text-align: center; margin-top: 10px;">18 MAY 1993</div>	Date of Mailing of this International Search Report <div style="text-align: center; margin-top: 10px;">17. 06. 93</div>													
International Searching Authority <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center; margin-top: 10px;">CHOULY J.</div>													

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

EP 9300599
SA 71427

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18/05/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4335125	15-06-82	US-A- 4223036	16-09-80
		US-A- 4144346	13-03-79
		AT-B- 366683	26-04-82
		AU-B- 521329	25-03-82
		AU-A- 3285078	09-08-79
		BE-A- 863382	27-07-78
		CA-A- 1094559	27-01-81
		DE-A- 2804096	03-08-78
		FR-A, B 2378778	25-08-78
		GB-A- 1594859	05-08-81
		JP-C- 1448813	11-07-88
		JP-A- 53095973	22-08-78
		JP-B- 62057634	02-12-87
		LU-A- 78960	21-06-78
		NL-A- 7801048	02-08-78
		SE-A- 7801088	01-08-78
		US-A- 4358449	09-11-82
		CH-A- 644118	13-07-84
		SE-B- 439773	01-07-85

WO-A-9102515	07-03-91	DE-A- 3927460	21-02-91
		AU-A- 6070790	03-04-91

EP-A-0490583	17-06-92	AU-A- 8837891	11-06-92
